

ANIMAL RESEARCH PAPER

Dairy farming: indoor v. pasture-based feeding

P. HOFSTETTER^{1*}, H.-J. FREY¹, C. GAZZARIN², U. WYSS³ AND P. KUNZ⁴

¹ Vocational Education and Training Centre for Nature and Nutrition (BBZN) Schüpfheim and Hohenrain, Chlosterbüel 28, CH-6170 Schüpfheim, Switzerland

² Agroscope, Institute for Sustainability Sciences (ISS), Tänikon, Switzerland

³ Agroscope, Institute for Livestock Sciences (ILS), Posieux, Switzerland

⁴ School of Agricultural, Forest and Food Sciences (HAFL), Bern University of Applied Sciences, Zollikofen, Switzerland

(Received 22 June 2013; revised 6 January 2014; accepted 18 February 2014; first published online 8 April 2014)

SUMMARY

The current situation of volatile milk prices and rising costs of, e.g. grain and labour, suggests that it is worth studying productivity and efficiency in dairy farming. The objective of the current whole-system study, carried out in lowland Central Switzerland from 2007 to 2010, was to compare the performance, efficiency, land productivity and profitability of indoor-feeding (IF) dairy production with that of pasture-based feeding (PF) dairy production. An IF herd consisting of 11 Holstein–Friesian (HF) and 13 Brown Swiss (BS) cows was kept in a free-stall barn and fed a part-mixed ration (PMR) of maize silage, grass silage and protein concentrate. The cows were allocated 15.8 ha of agricultural land (AL). In the PMR, an average per lactation of 443 kg protein concentrate and 651 kg compound feed was fed by a concentrate dispenser according to the requirements of each cow. The PF herd comprised 14 Swiss Fleckvieh (SF) and 14 BS cows, which were kept in a free-stall barn throughout the winter; barn-ventilated hay was offered *ad libitum* during the lactation period. This herd was allocated 15.7 ha of AL. After calving in spring, the PF cows grazed on semi-continuous pastures; they consumed an average of 285 kg of concentrate per lactation. The IF cows of the BS breed produced significantly more energy-corrected milk (ECM) per standard lactation compared with PF cows (8750 v. 5610 kg), more milk fat (350 v. 213 kg) and more milk protein (306 v. 203 kg). However, the milk of PF cows had higher levels of conjugated linoleic acid (CLA) (1.9 v. 0.6 g/100 g fat) and $\omega-3$ fatty acids (1.7 v. 0.9 g/100 g fat) than the milk of the IF cows. The calving interval (378 v. 405 days) and the empty time (87 v. 118 days) of the BS breed were significantly shorter in the PF in comparison with that of the IF production system. The IF herd yielded significantly higher ECM/ha AL and year (12 716 v. 10 307 kg), and showed a higher feed efficiency (1.3 v. 1.1 kg ECM/kg of total dry matter intake (DMI)). The productivity per hour was roughly similar in the two systems (IF: 76 v. PF: 73 kg milk/h). The PF system resulted in higher labour income compared with the IF system (20.7 v. 13.4 €/h), but the difference was not significant. In conclusion, land productivity and efficiency were higher with the IF herd than the PF herd due to the higher energy intake per kg feed. However, within the given conditions, the more interesting case, economically, might be the reduced costs and improved milk quality of the PF system rather than the increased milk yield of the IF cows.

INTRODUCTION

Recently, productivity and efficiency in dairy farming have gained in importance due to the increased cost of grain, following agricultural liberalization, and the rising cost of labour, machinery and buildings (Dillon *et al.* 2005; Macdonald *et al.* 2008). As a consequence

of the great demand for food (FAO 2013) and the limited availability of agricultural land, interest in improving the efficiency of small-scale farms has intensified. Efficient use of farm resources such as forage is becoming increasingly important.

Dairy farming in Central Europe is mainly based on indoor-feeding (IF) due to the long dormancy period of vegetation in the winter. During the vegetation period, cows are kept on half-day pasture and partly fed green

* To whom all correspondence should be addressed. Email: pius.hofstetter@edulu.ch

forage in the barn. In contrast, countries such as Ireland and New Zealand have adopted pasture-based feeding (PF) systems (Horan *et al.* 2005; Macdonald *et al.* 2008), which are characterized by low direct and overhead costs. Pasture-based feeding systems (FSs) now also play an increasingly prominent role in pre-Alpine regions. Comprehensive research has recently investigated the suitability of different breeds of dairy cows in different production systems, such as PF, IF with total mixed rations (TMR) or IF with part-mixed ration (PMR) (Holden *et al.* 1994; Horan *et al.* 2005; Burren *et al.* 2010; Horn *et al.* 2013). These studies have shown that the use of low-cost PF systems seems decisive for the survival of small-scale dairy farms in a highly competitive environment, especially when considering the rising cost of concentrate feed (Tozer *et al.* 2003; Gazzarin & Lips 2007; Alqaisi *et al.* 2011).

Notably, cows in PF production systems have lower milk solids content (Bargo *et al.* 2002; Vibart *et al.* 2008). Lower feed efficiency has also been reported in PF compared with IF cows due to limited energy intake on the pasture (Kolver & Muller 1998). Comparison of the studies by Bargo *et al.* (2002) and Fontaneli *et al.* (2005) confirms higher feed efficiency in IF systems.

To our knowledge, the existing literature lacks a whole-farm perspective on performance, efficiency, land productivity and labour income of the PF and IF systems that also considers the high European standards of environmental and animal welfare legislation. This gap is addressed in the current 3-year, full-lactation study.

On the basis of previous studies (Bargo *et al.* 2002; Fontaneli *et al.* 2005; Vibart *et al.* 2008), the following hypotheses were tested: (i) IF dairy cows are more efficient than PF dairy cows; (ii) based on a previous study (Münger & Jans 2001; Hofstetter *et al.* 2011), the land productivity of the herds would be similar; and (iii) the labour income of the PF herd would be higher (Durgjai 1996; Gazzarin & Schick 2004; Gazzarin *et al.* 2005).

MATERIALS AND METHODS

Experimental site

The experiment was performed on a mixed farm belonging to the Vocational Education and Training Centre for Nature and Nutrition in Hohenrain, Lucerne, which is in the lowland region of Central

Switzerland at 620 m above sea level. Annual precipitation was 1171 mm (SD = 131 mm) and the mean air temperature 9.4 °C (SD = 0.5 °C). In the vegetation period, the average precipitation was 946 mm and the mean air temperature 13.3 °C. Soils are medium-heavy, poor in humus, partly waterlogged, with sandy clay and brown earth. Dairy farming is the particular characteristic of land use in this region.

The botanical composition of the meadows was analysed in June 2010 using the method of Daget & Poissonet (1969). The swards of the IF systems, including the natural meadows, were on average composed of grasses (0.58), legumes (0.31) and herbs (0.11). On average, the pastures of the PF system consisted of 0.69 grasses, 0.22 legumes (only white clover, *Trifolium repens*) and 0.09 herbs. In the meadows, frequent grasses were perennial ryegrass (*Lolium perenne*) (0.31), smooth-stalked meadow grass (*Poa pratensis*) (0.22), rough meadow grass (*Poa trivialis*) and creeping bentgrass (*Agrostis stolonifera*). The forage growth period lasted from 10 March to 20 November.

Manure and chemical fertilizers were applied according to Swiss guidelines (Flisch *et al.* 2009), depending on the yield ability of the plots. On average, over three seasons, 177 kg N/ha, 128 kg P₂O₅/ha and 478 kg K₂O/ha/year were applied to the IF agricultural land (AL) and 166 kg N/ha, 104 kg P₂O₅/ha and 434 kg K₂O/ha/year to the PF AL.

Experimental design

The study was conducted from 1 October 2007 to 31 December 2010. Two systems, IF with mixed ration and PF, were examined over 3 years and 3 months, which covered three full lactations. Each system was allocated c. 16 ha of AL (Table 1).

The IF herd consisted of 24 dairy cows fed on PMR; part of the concentrate was distributed separately from a concentrate dispenser. In contrast to a TMR FS, in which cows are offered all forages as a complete ratio mix *ad libitum* (Linn 2013), PMR cows are fed a mixed basic ration, which is individually complemented by concentrate according to the requirements of each cow. In this herd, the target level was 8500 kg milk per lactation and cow in accordance with typical performance in local IF systems.

For PF, a seasonal, spring-calving, low-input, semi-continuous, pasture-based production system (Durgjai 1996; Münger & Jans 2001) with 28 dairy cows was established. In the PF herd, the target level was 6000 kg milk per lactation and cow, determined in

Table 1. *Agricultural land (AL) for the main forage and fodder crops for the energy and protein concentrate in hectares for the indoor-fed (IF) and pasture-fed (PF) herds, average from 2008 to 2010*

Area	IF		PF	
	Mean	SD	Mean	SD
Total AL (cropland included) (ha)	15.8	0.37	15.7	0.70
Main forage area (basic ration) (ha)	11.5	0.56	14.6	0.58
Pasture/hay (enclosed) (ha)	0.9	0.40	13.7	0.58
Grass silage (ha)	6.8	0.10		
Maize silage (ha)*	2.9	0.23		
Extensive meadow (ha) [†]	0.9	0.00	0.9	0.00
Fodder wheat meal (ha)*	0.8	0.19	0.5	0.11
Fodder maize meal (ha)*	0.6	0.15	0.4	0.16
Soyabean meal extract (or cake) (ha) [‡]	1.7	0.60	0.1	0.18
Maize gluten feed (ha) [§]	1.2	0.33	0.05	0.09

* In 2008 the majority of the maize silage, fodder wheat and fodder maize meal were purchased.

[†] Ecological compensating area.

[‡] The allocation for soyabean cake resulted from soyabean meal extract by an allocation of 67% and an output of 35.4 kg air-dried matter soyabean meal extract per ha.

[§] The allocation was 6.4% and by output of 71.6 kg air-dried material maize per ha.

accordance with previous pasture-based production system studies (Thomet *et al.* 2010; Hofstetter *et al.* 2011). At the beginning of each lactation, the PF cows were fed hay *ad libitum*, supplemented by limited amounts of concentrate.

The IF cows were housed in a free-stall barn during the whole lactation period, whereas the PF cows were kept in a free-stall barn during winter and the half-time grazing periods (early spring, late autumn). During the dry period, all cows were housed in a separate tied-stall barn.

Animals and animal measurements

Both herds included two breeds chosen to reflect the breeds generally used in corresponding FSs in Switzerland. As a result, herd composition partly differed between the two production systems.

On average, the IF herd consisted of 12.8 (SD=0.2) BS cows and 11.3 (SD=0.8) Swiss HF cows. The BS cows were 70 months old, had 3.6 lactations and a live weight (LW) of 697 kg. The HF cows were on average

64 months, had 3.1 lactations and an average LW of 700 kg. According to the herd-book data of the Swiss Brown Cattle Breeders' Federation (Braunvieh Schweiz 2012), the total merit index (TMI), i.e. the complete breeding value, of the BS was 108. The TMI (ISEL, i.e. complete breeding value with a combination of different breed indexes, weighted according to its economic importance) of the HF cows was 1033 according to the herd-book data of the Swiss Holstein Federation (Holstein Switzerland 2014). The IF cows were artificially inseminated with semen from sires of the same breed over the whole year.

The PF herd contained 13.8 (SD=0.3) BS cows and 14.2 (SD=0.7) Swiss Fleckvieh (SF) cows. The BS cows were 55 months old, had 2.6 lactations and a LW of 607, whereas the SF cows were 52 months, had 2.6 lactations and a LW of 617 kg. The TMI was 97 for the BS cows and 98 for the SF cows according to the cooperative Swiss Herdbook Zollikofen, Switzerland (Swiss Herdbook 2013). All TMI are based on April 2011 calculations. Because the merit ratings of the BS IF and PF cows were not calculated separately, those for the BS IF cows were slightly overestimated and those for BS PF cows slightly underestimated. The insemination period for PF cows lasted from 20 April to 20 July: they were artificially inseminated until the end of May, and afterwards mated by an Angus bull. They always calved between February and mid-April.

Cows were replaced as necessary, mainly as a result of infertility. Five IF cows were replaced in 2008, four in 2009 and eight in 2010 due to reduced fertility (five), mastitis (six), limping (two) and other causes (four). Four PF cows were replaced in 2008 due to fertility problems and a further one due to bloat; six were replaced in 2009, three of which were due to infertility; another six were replaced in 2010, four of which were due to infertility. On average, there were five primiparous cows in the IF herd and six in the PF herd.

Reproduction events and parameters (e.g. post-partum (p.p.) days open, empty time, services per conception, pregnancy and calving and calf's birth weight) were recorded. From calving to pregnancy, the cows were examined gynaecologically by an experienced veterinarian every second week to assess the status of the reproductive tract function and detect possible health problems.

Every 4 weeks, all cows were weighed at 6.30 AM, CET/CEST, after milking using EC 2000, modified Tru-Test scales with a precision of 1% (Tru-Test Ltd, Auckland, New Zealand). Body condition score (BCS)

was assessed every 2 weeks by the same scientist. The method of Edmonson *et al.* (1989), using a scale of 1–5, was applied in a slightly modified form (Metzner *et al.* 1993). Net energy for maintenance (MER) per day was calculated at $0.293 \text{ MJ net energy for lactation (NEL)} \times \text{metabolic live weight (i.e. LW kg}^{0.75}\text{)}$, according to Agroscope Liebefeld-Posieux (ALP) (2008). The same MER calculation was conducted for both production systems.

Feeding and grazing systems

During lactation, the IF cows were fed a PMR consisting of maize and grass silage, and protein concentrate. The basic PMR, formulated for 27 kg milk/day, was distributed from a chopping mixer wagon (LUCLEAR International) daily in summer, and every second day in winter. For a daily yield higher than 27 kg of milk, the IF cows were additionally fed a compound feed (CF), balanced according to individual requirements, from a concentrate dispenser (Lemmer Fullwood AG, Meierskappel, Switzerland). The ration and the amount of concentrate were calculated using the dairy ration analyser, a dairy feeding programme (CPM-Dairy, WexTech Systems Inc. 2006), and the nutrient requirements of dairy cattle (NRC 2001). During the vegetation period, IF cows grazed a siesta pasture for approximately 3 h/day. In the dry period, the IF cows were fed non-ventilated hay from the ecological compensation area and PMR leftovers in a tied-stall barn.

The PF cows were offered *ad libitum* access to barn-ventilated hay harvested from the surplus of the semi-continuous pasture. The amount of concentrate was limited to 4 kg concentrate per day at the beginning of the lactation (January–March). During the first month of the full-time grazing period, the PF cows were offered a maximum of 2 kg concentrate per day, supplemented by 4% magnesium oxide and mineral licks. The annual energy and protein concentrate limit was 8500 kg per PF herd. From March to mid-April and from mid-October to November, the cows spent half-days on the pasture and were additionally fed hay in the barn. From mid-April to mid-October, the cows grazed full-time except when they left the paddocks for c. 4 h/day for milking in the barn. In accordance with guidelines for pasture management from the Swiss Society for the Promotion of Fodder Production (Thomet *et al.* 1999), the PF herd grazed in a semi-continuous system consisting of four paddocks. Hay was harvested for winter feed from each paddock at

least once a year. During the dry period, the cows were fed non-ventilated hay from the ecological compensation area and straw in a tied-stall barn.

Forage storing, sampling and analysis

In accordance with the feeding programme based on milk performance and feeding requirements, the AL was compartmentalized into the required area for the different forages taking into account previous forage yields on the farm. Protein concentrate bought off the farm, i.e. soyabean or maize gluten feed, was converted into AL (Table 1) according to a study by Zimmermann (2006). Maize and grass silage were stored in round and cube bales. Hay was stored in the barn and partly ventilated.

The pasture yields were measured using the method proposed by Corral & Fenlon (1978), modified by Mosimann (2001). It was measured at four different and separate sites (10 × 1.0 m), at the border of the four paddocks (14 daily measurements, N fertilizer input of 200 kg and P fertilizer input of 100 kg/ha and year). Forage yields were determined from the average of the four sites at the border of the four paddocks.

When harvesting grass silage and hay, five to seven samples of 0.25 m² from each plot were cut at c. 60–70 mm above ground level with grass shears on the respective area to measure yields. The harvested yields were weighed using a Kern DE30K10 (Kern & Sohn GmbH, Balingen, Germany).

Every 2 weeks, pasture-grass samples were collected for nutritional analysis. The samples were cut at 30 mm above ground level on the stocked pastures with electric lawn scissors every six steps along the diagonal axis of the paddocks. Representative samples from the silage bales were collected 1 month before feeding. Maize silage samples were collected once in November, 6 weeks after harvesting. Each year, a mixed hay sample from each haystack was obtained by thrusting a lance diagonally in three different places.

The pasture forage was sampled from the beginning of April until the beginning of November. The collected forage (c. 1.5 kg of fresh pasture) was dried to a constant weight, one half at 55 °C and the other half at 105 °C in a forced-air oven for 24 h to determine dry matter (DM) content. Pasture-forage samples and all other forage samples were analysed at the forage testing laboratory of Dairy One (Ithaca, New York) in accordance with the official methods of analysis (AOAC 2005) method 989.03 using Near-Infrared Reflectance Spectroscopy (NIRS) (Foss NIR Systems

Model 6500 with ISI II v1.5), following Van Soest *et al.* (1991).

Measurement of milk performance and milk solids

Procedure A4 of the Braunvieh Schweiz (2013) standard milk yield recording protocol and guidelines was followed to measure individual milk yield in the morning and in the evening. Measurements were conducted twice per month, using the Afimilk 2000 milk meter (Boutech AG, Baar, Switzerland). From 1 October to 31 December 2007, six measurements of the IF cows were taken. Twenty-two measurements of each were taken in 2008 and 2009 and 23 values were obtained for 2010. The lactation persistence was calculated as the percentage of the milk performance of the second 100 days in relation to the first 100 days milk performance. Analyses of fat, protein and lactose concentration of the BS and SF cows were performed at Qualitas AG Laboratory (Zug, Switzerland), in accordance with the recording protocol. Analyses of fat, protein and lactose concentration of the HF cows were performed at the laboratory of the Swiss HF breeders' association in Posieux, Switzerland. The cows were milked at 5:15 AM and at 4 PM, CET/CEST, in a 2×5 herringbone milking parlour (Lemmer Fullwood AG, Meierskappel, Switzerland). Every month, tank milk samples of the two systems were taken and the fatty acid (FA) composition of the milk fat was analysed according to Collomb & Bühler (2000). Energy-corrected milk (ECM) was calculated according to Sjaunja *et al.* (1990) as $(0.038 \times \text{g fat/kg milk} + 0.024 \times \text{g crude protein/kg milk} + 0.017 \times \text{g lactose/kg milk}) \times \text{kg milk}/3.14$.

Feed intake, feed efficiency and productivity

The feed efficiency for both systems was calculated as kg ECM produced per kg of total dry matter intake (DMI). The net energy for lactation (in MJ NEL) per herd was calculated by multiplying the weighted yield of the feed by the corresponding energy value. For every cow and herd, the required daily and the consequent annual net energy was calculated by adding the kg of ECM $\times 3.14$ MJ NEL and the maintenance energy requirement. Energy for calf growth during pregnancy was considered only for months 8 and 9, adding 11 and 18 MJ/day, respectively. Energy for the cow's growth was not considered.

The productivity of the AL was determined by dividing the annual ECM production per herd by the

available area, including the ecological compensation area, in accordance with AGFF guidelines (AGFF 2002). The annual ECM from the basic ration was determined by the AL productivity minus the milk potential from concentrate using calculations as shown below.

Milk production potential (MPP) of a kg concentrate or forage (in DM) = corresponding MJ NEL/3.14 MJ NEL

ECM from basic ration (kg/ha) = ECM (kg/ha AL) – MPP total concentrate/ha AL

ECM from basic ration (kg/cow) = ECM (kg/cow) – MPP concentrate/cow

Cost and profitability

Accounting for the full cost analysis was done separately for the two systems and the two herds. All incomes, direct costs and most of the overhead costs were allocated to the respective accounts, which were audited by the agricultural fiduciary Agro-Treuhand Sursee (Sursee, Switzerland). Full cost analysis was conducted using International Farm Comparison Network (IFCN) methods.

The variable costs of the concentrate feed for the IF and PF systems were calculated by subtracting the proceeds of crop (wheat and maize meal) sales from costs of purchased concentrates. The net variable costs were subsequently debited against the IF account.

The fixed costs were allocated using standard costs from different calculation programmes (Gazzarin & Hilty 2002; Gazzarin & Schick 2004; Hilty *et al.* 2007). Machinery costs were allocated according to the time the tractors were used in each system, namely 302 h (0.76) for IF and 98 h (0.24) for PF.

The calculation of work time was done by combining measured work time in the present study with standard data from a previous analysis of similar production systems (Stark *et al.* 2009). The total AL in both systems was notionally divided into one-half rented and one-half owned areas. Rent for the AL was set at 666 €/ha and year, using standard values. Furthermore, the notional rent of owned land was calculated as opportunity costs in fixed costs. Overall, 0.8 of the total operating costs (e.g. telephone, insurance) were allocated to both systems.

Statistical analysis

Data were recorded from 1 October 2007 to 31 December 2010. Information about forage




Fig. 1. Course of the average Body Condition Scores (mean and SD) of the indoor and the pasture-fed cows during lactation in 2007–2010. Average measurements per period (n) were Brown Swiss (BS) IF 74, BS PF 85, Holstein–Friesian (HF) 64 and Swiss Fleckvieh (SF) 87.

production, the calculation of DMI and productivity as well as costs and revenues of the two systems was compiled on an annual basis. The data are given as means and SD for the 3 years.

The annual period considered for the cow data in the PF system was from 1 January to 31 December each year. For the IF system, the data collection period was from 1 October to 30 September, except for the final year, which lasted until 31 December; data collection started earlier for this system due to the longer calving intervals.

Data for individual cows were recorded per lactation. Consecutive lactations of the same cow were treated as independent data. This simplification was necessary because some of the cows were replaced over time and because lactations of the IF cows were not synchronized, i.e. data did not have a classical repeated-measures structure.

Each lactation was subdivided into time segments of 28 days, beginning p.p. Measurements for BCS and LW began 28 days prior to calving, and those for milk production began at calving. Measurements taken every 14 days were attributed to the 28-day segments for each cow according to calving date. For each time segment, moving averages from 14 days prior and up to 14 days after were calculated. Data from all time segments are represented graphically in Figs 1 and 2,

but only data from four time segments (29–56, 57–84, 85–112 and 281–308 days), as well as total values per lactation, were tested statistically.

The four groups of cows (two systems, each with two breeds) were compared using means and standard errors (S.E.) of the performances per lactation recorded from each group over the study period. However, the significance of the differences between the systems was tested using only data from BS cows. Fertility and milk variables were compared between the two FS and the years using two-way ANOVA, non-repeated-measures ANOVA for the reason given above.

The Pearson product–moment correlation was used to describe correlations between variables. All analyses were conducted using R-2.11.1 software by R Development Core Team (2012).

RESULTS

Forage yield and quality

The strongest average meadow growth of 105 kg DM/ha/day occurred at the beginning of May. The growth decreased at the end of May and in June. From July to mid-September, it increased again and reached 80 kg DM/ha/day before it decreased continuously throughout the autumn. The total yield, calculated according




Fig. 2. Course of the average milk yields (kg ECM, mean and SD) of the indoor and the pasture-fed cows during lactation in 2007–2010. Average measurements (n) per period were: Brown Swiss (BS) IF 59, BS PF 70, Holstein–Friesian (HF) 55 and Swiss Fleckvieh (SF) 72.

to Corral & Fenlon (1978), was 14.2 (SD=0.5) t DM/ha/year.

Table 2 shows the chemical composition and energy content of the basic ration and the concentrate of both systems from 2008 to 2010. The pasture had high crude protein (CP) content: from 2008 to 2010, the CP content of the pasture increased continuously (Table 3) with a significant difference between 2008 and 2010 ($P<0.05$). No significant differences were found between the years in terms of water-soluble carbohydrates (WSC), acid-detergent fibre (ADF) and NEL in the pasture. However, in 2009, the neutral-detergent fibre (NDF) content of the pasture was significantly lower compared with 2010 ($P<0.05$). A comparison between the spring and summer values showed significant differences in the dietary components of the pasture (ADF: $P<0.001$, NDF: $P<0.01$, WSC: $P<0.001$, NEL: $P<0.001$), except for CP (Table 3).

In 2008, 2009 and 2010, the energy content of grass silage was 6.1, 6.1 and 6.0 MJ NEL/kg DM, respectively, the average CP content was 164, 160 and 174 g/kg DM, the average ADF content was 312, 318 and 313 g/kg DM, and the average NDF content was 463, 491 and 470 g/kg DM.

The measured average annual yield of grass silage was 13.8 (SD=0.8) t DM/ha and for maize silage was 17.7 (SD=1.9) t DM/ha. The first cut comprised 4.8 t DM/ha, the second 2.7 t DM/ha, the third 2.7 t DM/ha, the fourth 2.0 t DM/ha and the fifth 1.6 t DM/ha of the grass silage.

The average annual yield of fodder wheat was 7.9 (SD=1.0) t air-dried matter/ha and of fodder maize 9.5 (SD=1.9) t air-dried matter/ha.

Feed intake, feed efficiency and productivity

The calculated energy intake in NEL of the IF herd was mainly based on grass silage (419 123 MJ), maize silage (334 448 MJ) and concentrate (202 357 MJ). Energy intake in NEL from pasture (46 360 MJ) and from hay from extensive meadows (32 785 MJ) was low. The calculated energy intake in NEL of the PF herd was based mainly on pasture (571 467 MJ) and hay (230 632 MJ); that from concentrate (63 048 MJ), hay from extensive meadows (37 797 MJ) and straw (6910 MJ) was low.

The annual DMI of the IF herd was 157.6 (SD=4.6) t and that of the PF herd 150.1 (SD=6.8) t. The total DMI of an IF cow, including the dry period, was

Table 2. Chemical composition and energy content of the basic ration and of concentrate offered during the study to the indoor-fed (IF) and pasture-fed (PF) herds. Dry matter (DM); crude protein (CP); neutral detergent fibre (NDF); acid detergent fibre (ADF); water-soluble carbohydrate (WSC); net energy for lactation (NEL)

System	Year	Feed	n	DM (g/kg)		CP (g/kg DM)		NDF (g/kg DM)		ADF (g/kg DM)		WSC (g/kg DM)		NEL (MJ/kg DM)	
				Mean	SD	Mean	Mean	SD	SD	Mean	SD	Mean	SD	Mean	SD
IF	2008–10	Grass silage	48	449	82.3	165	29.3	448	47.1	307	24.5	140	18.6	6.1	0.42
IF	2008–10	Maize silage	3	369	20.4	85	5.0	363	31.4	215	20.7	37	17.5	7.3	0.22
PF	2008–10	Pasture	51	159	27.0	260	27.1	379	41.2	242	25.6	132	10.6	6.3	0.42
PF	2008–10	Hay/aftermath	5	873	15.9	161	28.8	427	52.3	291	24.5	160	19.3	6.1	0.39
IF/PF	2008/09	Protein concentrate (PC)*	1	890		584		92		48		31		7.8	
IF	2010	Maize gluten feed	1	890		697		110		68		16		8.5	
PF	2008–10	Energy concentrate (EC) [†]	1	890		117		99		38		32		8.1	
IF	2008–10	Compound feed (CF) [‡]	1	890		198		143		57		28		8.5	

* 475 g/kg soyabean cake, 475 g/kg maize gluten feed, 35 g/kg mono-calcium phosphate and 15 g/kg calcium carbonate, salt and further minerals. In 2007, PF cows were fed PC.

† 487 g/kg maize meal, 487 g/kg wheat meal, 20 g/kg sugar beet molasses, 3 g/kg mono-calcium phosphate, 3 g/kg calcium carbonate.

‡ 358 g/kg maize and wheat meal, 157 g/kg soyabean cake, 77 g/kg maize gluten feed, 15 g/kg sugar beet molasses, 18 g/kg crystalline fat, 7 g/kg mono-calcium phosphate and 10 g/kg calcium carbonate.

Table 3. Chemical composition and energy content of pasture offered per year and in spring and summer. Crude protein (CP), acid detergent fibre (ADF), neutral detergent fibre (NDF), water-soluble carbohydrate (WSC), net energy for lactation (NEL).

	n	2008		2009		2010		P value year			April		August		P value	
		Mean	SD	Mean	SD	Mean	SD	2008 v. 2009	2008 v. 2010	2009 v. 2010	n	Mean	SD	Mean	SD	Season
CP (g/kg DM)	17	249	27.0	259	23.3	272	27.4	0.286	0.022	0.146	9	258	36.5	279	11.9	0.128
ADF (g/kg DM)	17	243	24.4	238	28.2	247	21.4	0.545	0.657	0.299	9	213	25.0	259	5.4	<0.001
NDF (g/kg DM)	17	385	40.9	364	40.8	392	35.1	0.148	0.597	0.041	9	343	45.7	404	11.6	0.006
WSC (g/kg DM)	17	143	50.6	131	33.1	122	30.3	0.418	0.148	0.405	9	183	43.5	86	7.8	<0.001
NEL (MJ/kg DM)	17	6.3	0.6	6.4	0.4	6.2	0.3	0.539	0.853	0.246	9	6.6	0.3	6.0	0.2	<0.001

Table 4. Concentrate offered, energy content of offered ration, production intensity, land productivity, feed efficiency and average calf birth weight of the indoor-fed (IF) and pasture-fed (PF) herds, average from 2008 to 2010

	IF		PF		P value
	Mean	SD	Mean	SD	
Concentrate (g air-dried/kg ECM)*	131	14.7	53	6.0	0.001
NEL in DMI (MJ/kg) [†]	6.6	0.02	6.1	0.10	0.001
Production intensity (MJ/MJ) [‡]	2.9	0.04	2.4	0.05	<0.001
ECM (kg/ha AL)	12716	201.3	10307	616.5	0.003
ECM from basic ration (kg/ha)	8808	321.0	9039	663.1	0.616
ECM from basic ration (kg/cow)	5699	278	5022	155	0.021
ECM/DMI (kg/kg), i.e. feed efficiency	1.3	0.04	1.1	0.03	0.002
ECM/kg of LW ^{0.75} (kg)	62	1.3	47	1.0	<0.001
Calves' birth weight (kg)	1076	29.3	1273	35.7	0.002
Calf's birth weight (kg/ha AL)	67	2.9	81	5.1	0.014

* Energy and protein concentrate without added mineral nutrients.

† DMI; dry matter intake.

‡ Production intensity: Net energy total (NEL_{total}; kg ECM ×3.14 MJ) plus net energy for maintenance (NEL_{maintenance}; kg metabolic LW ×0.293 MJ ×365)/NEL_{maintenance} (ALP 2008).

17.9 (SD=0.3) kg/day and of a PF cow 14.6 (SD=0.4) kg/day.

The IF herd had a higher-energy concentration in the feed and a higher production intensity compared with the PF herd (Table 4). The cows of the IF herd produced significantly more ECM per hectare AL and year ($P<0.01$) and significantly more ECM per kilogram of metabolic LW ($P<0.001$). The ECM from basic ration was higher in the PF system, but insignificant, whereas ECM from basic ration/cow was significantly higher in the IF system ($P<0.05$). The IF cows had a significantly higher feed efficiency (kg ECM/DMI) compared with the PF cows. The PF herd showed higher ECM performance per hectare and year from the basic ration, but this was insignificant. On average, the PF herd produced a significantly higher average calf birth weight kg/year ($P<0.01$) and kg/ha ($P<0.05$) than the IF herd.

Grazing period, stocking rate, live weight and body condition score

On average, the full-time grazing period of the PF herd lasted 179 days and total grazing period 242 days. The average stocking rate in the IF herd was higher than that of the PF herd, at 2.1 v. 1.9 cows/ha of main forage area.

Before calving, no significant between-system differences were found in the LW (data not shown) and

BCS (Fig. 1). However, in the segments at the beginning and end of lactation, the IF cows had significantly higher LW (segment 1: $P<0.05$, 2: $P<0.01$, 3: $P<0.01$ and 4: $P<0.01$). The lowest LW of the BS cows was observed at 58 (IF) v. 119 (PF) days p.p. ($P<0.001$), whereas that of BCS was seen at 92 (IF) v. 168 (PF) days p.p. ($P<0.001$), both significantly later in the course of lactation in the PF than in the IF cows. The LW loss of IF BS cows correlated negatively with milk protein content (Pearson's $r=-0.59$, $n=18$, $P<0.01$). Milk yield in the first segment of lactation correlated negatively with the lowest LW of BCS in the PF cows ($r=-0.64$, $n=28$, $P<0.001$), but was not significant for the IF cows of the BS breed. Milk yield in the second segment correlated negatively with the lowest LW of BCS in the PF cows ($r=-0.56$, $n=28$, $P<0.01$), but not in the IF cows.

Fertility

During the study, four of the IF cows calved in the first quarter of the year, 13 in the second, 34 in the third and 18 in the fourth. In contrast, seven of the PF cows calved in January, 47 in February, 25 in March and nine in April.

The PF cows had significantly shorter calving intervals, empty time and time from calving to first service (Table 5). The IF cows showed significantly shorter pregnancy duration ($P<0.05$), but the difference was

Table 5. Comparison of fertility dates between feeding systems and cow breeds. Mean and s.e. are given for each of the four cow groups

	Brown Swiss									Holstein–Friesian			Swiss Fleckvieh		
	Indoor feeding			Pasture feeding			P value			Indoor feeding			Pasture feeding		
	n	Mean	s.e.	n	Mean	s.e.	FS*	Year	FS × year	n	Mean	s.e.	n	Mean	s.e.
Services per conception	52	1.9	0.17	43	1.8	0.17	0.570	0.167	0.936	34	2.3	0.27	43	1.5	0.13
Calving interval (days)	29	405	9.5	33	378	5.8	0.021	0.684	0.423	26	406	13.2	41	370	4.2
Time empty (days)	31	118	8.9	34	87	5.6	0.003	0.363	0.379	26	125	13.1	41	83	4.3
Calving to first service (days)	31	82	2.8	34	69	3.0	0.022	0.023	0.173	26	87	5.1	41	71	3.2
First service to conception (days)	31	36	8.2	34	18	4.4	0.064	0.601	0.386	26	38	11.8	41	12	2.9
Pregnancy duration (days)	44	288	1.0	41	291	0.9	0.042	0.531	0.193	28	280	0.9	45	286	0.8
Calves' birth weight (kg)	45	44	0.9	43	44	0.8	0.728	0.046	0.982	28	45	0.9	45	43	0.9

* FS, feeding system.

Table 6. Comparison of milk production variables (milk yield and milk solids) between feeding systems and breeds in standard lactation. Mean and s.e. are given for each of the four cow groups

	Brown Swiss*							Holstein–Friesian*		Swiss Fleckvieh*	
	Indoor feeding		Pasture feeding		P value			Indoor feeding		Pasture feeding	
	Mean	s.e.	Mean	s.e.	FS†	Year	FS × year	Mean	s.e.	Mean	s.e.
Milk yield (kg/lactation)	8481	250	5839	180	<0.001	0.027	0.851	9348	302	6253	184
ECM (kg/lactation)	8750	259	5610	170	<0.001	0.042	0.518	9422	299	6082	164
Milk fat (g/kg)	41	0.4	36	0.6	<0.001	0.584	0.105	41	0.5	38	0.6
Milk fat (kg/lactation)	350	11.2	213	7.2	<0.001	0.082	0.334	380	13.1	240	6.8
Milk protein (g/kg)	36	0.3	35	0.3	<0.01	0.165	0.014	34	0.3	34	0.2
Milk protein (kg/lactation)	306	8.1	203	5.6	<0.001	0.017	0.556	317	9.3	209	5.7
Lactation persistence (%)	87	2.0	67	1.4	<0.001	<0.001	0.352	85	1.7	70	1.1

* n: BS IF; 32, BS PF; 29, HF; 30, SF; 38.

† FS, feeding system.

Table 7. Saturated, mono- and poly-unsaturated fatty acids (FA), conjugated linoleic acid (CLA) and $\omega-3$ FA of the cow milk fat from the feeding systems, average from 2008 to 2010

	Indoor feeding*		Pasture feeding*		P value		
	Mean	S.E.	Mean	S.E.	FS [†]	Year	FS × year
Saturated FA (g/100 g fat)	63.5	0.45	57.7	0.48	<0.001	0.014	0.527
Mono-unsaturated FA (g/100 g fat)	21.5	0.39	26.0	0.42	<0.001	0.338	0.202
Poly-unsaturated FA (g/100 g fat)	3.6	0.13	5.5	0.14	<0.001	0.165	0.386
CLA (g/100 g fat)	0.6	0.09	1.9	0.09	<0.001	0.078	0.393
$\omega-3$ FA (g/100 g fat)	0.9	0.03	1.7	0.04	<0.001	0.348	0.679

* *n*: for the IF system 12 each year; for the PF system 10 (2008), 11 (2009) and 11 (2010).

† FS, feeding system.

only 3 days. No significant differences were found in services per conception and time from first service to conception. Time from calving to first service was significantly different across the years ($P < 0.05$). Time from first service to conception correlated significantly with milk yield in the IF BS cows ($r = 0.53$, $n = 17$, $P < 0.01$), but not in the PF cows. There was no significant difference in the average calf birth weight between the IF (44.2, $SD = 5.5$ kg) and PF (43.4, $SD = 5.7$ kg) cows.

Milk performance and milk solids

Significant differences ($P < 0.001$) between the IF and PF cows of the BS breed were found in milk yield, milk fat, milk protein production (kg/lactation) and in lactation persistence in addition among the year (Table 6). Milk protein content is also different ($P < 0.01$) between the FS. A significant interaction occurred between FS and year in milk protein content (g/kg) ($P < 0.05$). The lactation curves of the average milk yields show the lactation persistence in particular (Fig. 2).

The milk solids (data not shown) indicate that the milk of the IF cows had a significantly higher fat content than that of the PF cows in segments 1, 2 and 3 ($P < 0.01$), but not at the end of lactation. The milk from the PF system had significantly fewer saturated and more mono- and poly-unsaturated FA, as well as more conjugated linoleic acid (CLA) and more $\omega-3$ FA than that of the IF system (Table 7). Insignificant differences in the average milk protein content were found at the beginning of lactation, in segment 1. Compared with PF, the IF cows showed a significantly higher milk protein content in segments 2 ($P < 0.05$) and 3 ($P < 0.01$), but a significantly lower protein content at the end of the lactation ($P < 0.01$).

The PF BS cows showed a significant increase in milk urea content to a peak of 54.9 (s.e. = 1.3) mg/dl on the 170th day of lactation (segment 1: $P < 0.01$, segment 2: $P < 0.01$), whereas the IF cows showed no significant fluctuation in the average of milk urea content, with a range of 25.0–28.4 mg/dl over the course of lactation. At the end of lactation, the urea content of the milk was similar in both herds. The milk sugar content of the IF BS cows decreased continuously from 48 g/l in the first segment to 45 g/l in the fourth. Compared with PF BS cows, the IF BS showed a significantly higher average milk sugar content at the end of lactation ($P < 0.01$) and between days 57 and 84 of lactation ($P < 0.05$).

Cost and profitability

Significantly higher milk production was obtained from the IF herd than from the PF herd ($P < 0.01$), but there was no significant difference in milk returns (€/100 kg). In contrast, the calves' returns were significantly higher from the PF system compared with the IF system ($P < 0.05$, Table 8). Direct costs of concentrate ($P < 0.05$), hired labour and machinery costs (contract labour included) ($P < 0.001$) were, however significantly lower for the PF herd. Neither equipment nor building costs reached statistical significant differences among the FS. Although the PF system had significantly higher operating costs ($P < 0.001$), its labour income per hour still tended to be higher; this result, however, did not reach statistical significance ($P < 0.10$).

DISCUSSION

The principal aim of the current full-lactation study was to investigate the performance, land productivity

Table 8. Performance, direct and fixed costs, labour income and productivity of the indoor-feeding (IF) and pasture-based feeding (PF) systems, average from 2008 to 2010

	IF		PF		P value
	Mean	SD	Mean	SD	
Produced milk (t/year)	194	5.0	165	4.3	0.003
Milk returns (€/100 kg)*	53.5	8.86	50.6	8.48	0.706
Cull cow returns (€/100 kg)	4.2	1.68	3.3	0.56	0.419
Calves (€/100 kg)	3.9	1.14	6.6	1.06	0.043
Direct payments [†] (€/100 kg)	14.0	1.45	16.8	1.35	0.075
<i>Direct costs</i>					
Animal purchases (€/100 kg)	8.7	2.75	9.3	0.75	0.750
Purchased feed [‡] (€/100 kg)	7.7	1.61	3.9	0.59	0.019
Veterinarian [§] , insemination (€/100 kg)	6.0	0.44	5.5	0.35	0.207
Miscellaneous animal costs (€/100 kg)	2.0	0.36	2.4	0.33	0.285
Forage crop husbandry (€/100 kg)	3.3	0.13	3.1	1.03	0.854
<i>Fixed costs</i>					
Contract labour, hire machinery (€/100 kg)	11.5	1.40	1.1	0.35	<0.001
Own machinery (€/100 kg)	6.1	1.30	4.7	0.63	0.173
Buildings, interest included (€/100 kg)	9.0		11.8		
General operating expense (€/100 kg)	3.2	0.28	6.3	0.33	<0.001
<i>Income</i>					
Agricultural income (€/100 kg)	18.2	5.54	29.1	6.04	0.082
Opportunity costs of own capital (€/100 kg)	0.6		0.7		
Labour income (€/100 kg)	17.6	4.6	28.5	4.2	0.083
<i>Productivity</i>					
Total manpower requirements (h/year)	2553		2268		
Labour productivity per labour unit (kg milk/h)	76	2.0	73	1.9	0.178
Labour income per labour unit (€/h)	13.4	3.48	20.7	3.07	0.092

* Average milk price/100 kg in 2008, 2009 and 2010 was 67.0, 52.0 and 51.2 € for the IF system and 64.5, 48.3 and 48.7 € for PF, respectively. Currency exchange rate: 1 CHF=0.8325€.

† Including ecological payments.

‡ Considered the inter-farm supply for cereals.

§ Monitoring of health included.

|| Own fodder crops, i.e. seed, fertilizer.

and the agricultural and labour income of PF and IF dairy herds in a region with a limited vegetation period. This is the first such study to investigate differences in productivity and efficiency between BS, SF and HF cows in two different systems under equal conditions. The current market situation, with its risk of rising feed prices due to greater demand (FAO 2013), agricultural reforms, drought and ecological considerations may indicate that self-sufficient farming using local resources (e.g. pasture) on smallholder farms will become more important in the future.

Forage yield and quality

The average grass yields of the systems were typical for the Swiss lowlands (Thomet & Blättler 1998;

Hofstetter *et al.* 2011). Sufficient grass yield can be achieved in swards mixed with grass, herbs and legumes using moderate levels of N fertilizer (about 170 kg N/ha/year) according to Nyfeler *et al.* (2011). Nitrogen application in the present study was lower than that used in Moorpark, Ireland (McEvoy *et al.* 2008, 2009; Curran *et al.* 2010), but similar to that used in New South Wales, Australia (Fariña *et al.* 2011) and in Florida, USA (Fontaneli *et al.* 2005).

Compared with the average Swiss values (ALP 2008; Schori 2009), maize silage and hay were higher in energy content, in NDF and ADF values, and grass had higher CP concentration. Crude protein in pasture was similar or higher compared to that reported by McEvoy *et al.* (2009) and Curran *et al.* (2010). Bargo *et al.* (2002) reported higher values in pasture; Vibart *et al.*

(2008) found lower CP and ADF values but higher NDF values. Differences in CP concentration may be due to grass composition, pasture management and N fertilization. Furthermore, differences in ADF, NDF and NEL may be explained by minor differences in the method of chemical analysis and different regression calculations for NEL. High CP in meadows and hay is partly explained by its high leguminous content (Nyfeler *et al.* 2011). A potential explanation for increased CP concentration in pasture from 2008 to 2010 might be a corresponding increase in total harvested N over the 3 years, an improvement in pasture management, or adjustment of the swards to the PF systems.

The variation in chemical composition and energy content of pasture within the vegetation time, e.g. spring *v.* summer, was higher than the variation in nutrient values over the 3 years; similar data were shown by Fariña *et al.* (2011). Variation in grass content has an important impact on milk yields. No significant between-year differences were found, except for CP and NDF. As a result of this near-balanced data set, the PF herd data from different years has been pooled for analysis. The between-year variation of grass silage was smaller than the seasonal variation and the data set of the IF systems has also been pooled.

Feed intake, feed efficiency and productivity

The higher energy intake and the significantly higher production intensity in the IF herd resulted in higher feed conversion efficiency (kg ECM/kg DMI) and in more ECM/kg of metabolic LW. In contrast, with less feed intake, lower milk performances and lower production intensity the PF cows required a higher share of their energy intake for their MER. A higher energy requirement in the PF cows due to the physical activity on the pasture might have resulted in a lower coefficient of milk efficiency (Kaufmann *et al.* 2011).

The feed conversion efficiency in the PF cows was slightly lower than the results of Schori & Münger (2010). Their average milk productivity per metabolic LW was consistent with previous studies (Steiger Burgos *et al.* 2007; Schori & Münger 2010; Thomet *et al.* 2010). The feed conversion efficiency of the PF herd was also lower compared to the results of Fariña *et al.* (2011), who used more concentrate, a higher stocking rate and different ECM calculation methods. The clearly higher milk production of the IF

cows led to the higher milk performance from the basic ration.

Productivity (kg ECM/ha AL) was high in the IF herd compared with the PF due to the higher energy and protein content of the feed ration, the higher feed intake and the higher milk production. The productivity of the PF herd was lower than the results reported by Hofstetter *et al.* (2011), who used a rotational grazing system, and also lower than the findings of both McEvoy *et al.* (2009) and Curran *et al.* (2010). Their swards, however, were already adapted to the PF system, whereas the swards in the present study were used as semi-continuous pastureland for the first time. However, McEvoy *et al.* (2009) and Curran *et al.* (2010) used a stocking rate twice as large, did not analyse the whole lactation, and worked in more favourable climatic conditions.

The PF herd did not show a significantly higher productivity from the basic ration than the IF herd, but the performance/cow from basic ration was significantly higher ($P < 0.05$) for the IF cows compared with the PF ones. The less-than-outstanding land productivity for the PF herd is probably related to the feed intake during the summer. It seems that the PF cows could not eat enough forage from the short swards offered in the experiment. This is in contrast to the comparative studies between continuous pasture and rotational grazing in Kent (UK) by Pulido & Leaver (2003) and in Switzerland by Münger & Jans (2001), who fed more concentrate, and in France by Hoden *et al.* (1987), who applied more N fertilizer. Further studies should investigate whether additional feed and the use of a rotational grazing system or other measures would improve the economic results of pasture land FSs.

Stocking rate, live weight and body condition score

In the present study, the stocking rate was defined as cows per hectare because it is based both on equal conditions regarding AL and cows suitable for the respective systems. However, if the appropriate stocking rate is considered to be 600 kg LW/ha, the calculated stocking rate would be 2.4 cows/ha for the IF farm and 2.0 cows/ha for the PF farm.

The LW loss in the IF cows corresponds with the findings of Fontaneli *et al.* (2005), but the PF cows in Fontaneli *et al.* (2005) showed 30 kg higher LW losses than our PF cows. The long-lasting LW loss of the PF cows during the summer probably resulted from lower seasonal yields, despite the fact that the stocking

rate was consequently reduced, e.g. in 2010 from 4.1 cows/ha in April to 2.8 cows/ha in mid-June. In June and August, the pasture had a lower WSC concentration and lower energy content. Thus, energy intake and grass consumption were lower, in agreement with the results of Kolver & Muller (1998). The low energy intake resulted in a relatively long-lasting energy deficiency over the summer. This might explain the lower fat and protein content in the milk of the PF cows at the beginning of lactation and until the 150th day of lactation.

Despite an absence of significant differences in daily LW loss among FS, a negative correlation between daily LW losses and milk protein content in the IF herd suggests a relationship between LW loss and protein content in the milk. No such correlation was found in the PF herd, which showed no pronounced decrease in daily LW losses over a short time. The significant negative correlation between milk performance in days 1–56 p.p. and the lowest LW for BCS suggests insufficient energy supply over the summer in the PF cows. No such correlation was found in the IF cows.

Fertility

Services per conception, especially in the IF cows, were high compared to results reported by Reist *et al.* (2000). The empty time of IF BS cows was 2 days shorter than the average for Swiss dairy cows in 2010/11 (Braunvieh Schweiz 2011). Compared with PF cows, IF cows showed significantly longer empty time and a tendency towards a longer time from first service to conception, but the differences between the systems were not significant. These findings are consistent with the studies of Vibart *et al.* (2012). A positive correlation between time from first service to conception and milk yield illustrates that dairy cows with high milk performance are disadvantaged with more services.

In the current PF system, the cows showed a higher number of services per conception, fewer days in the calving interval and less time empty compared to the results from a low-input production system in an Austrian Alpine region with Brown Swiss (BS) and adapted Holstein–Friesian (HF) cows (Horn *et al.* 2013). This difference occurred even though these cows were fed more concentrate supplementation per year. According to the studies by Butler & Smith (1989), Barb & Kraeling (2004) and Butler (2005), a high daily LW loss in a short period of time until reaching the lowest point, as found in the IF BS cows

(–1228 g/d), may result in longer empty time. The longer calving interval and longer empty time of the IF system is probably due to the temporally defined insemination period and the presence of a bull on the pasture in the PF system.

Due to the PF system's higher number of cows and a standard LW of 600 kg/cow, the PF herd produced higher average calf birth weight per AL compared with the IF herd. Fertility in the present study is similar to results from previous studies (Coleman *et al.* 2009; Vance *et al.* 2012). More cows might be needed to provide stronger evidence regarding reproductive performance.

Milk performance and milk solids

The different lactation curves for the IF and PF cows are consistent with the results of Fontaneli *et al.* (2005) and Bargo *et al.* (2002) and in accordance with the PF system of Fariña *et al.* (2011). In contrast to the study of Bargo *et al.* (2002), the differences in milk production during the whole lactation were significant in the present study. The IF cows showed a stable course of the average milk urea content, which suggests that the feed ration was appropriate to requirements. In contrast, PF cows showed the typical peak of milk urea. Low WSC and low energy content in the grass during the summer partly explains the lower milk-fat-to-milk-protein ratio in PF cows compared with IF cows. The results showing higher milk performance and milk solids in the IF cows compared with the PF cows are broadly consistent with previous comparative studies (Fontaneli *et al.* 2005; Vibart *et al.* 2008, 2012).

The higher content of CLA and $\omega-3$ FA of the PF cows confirm the results of Dhiman *et al.* (1999), White *et al.* (2001) and Collomb *et al.* (2008), regarding the positive influence of mountain pasture on milk quality in Switzerland. In particular, from April to October the PF cows showed fewer saturated but more unsaturated fatty acids, and more CLA and $\omega-3$ FA compared with the cows fed on conserved forage (Wyss *et al.* 2011).

The PF cows showed a constant increase in milk performance of about +200 kg ECM/cow/year and an increase in milk protein. This might be a result of an adaptation process of metabolism to the PF system and is consistent with results from a previous study (Hofstetter *et al.* 2011). However, it might also have been a consequence of improved management. The IF cows did not show this increase in milk performance in the study period.

The suboptimal feed and energy availability in the continuous pasture-FS, especially during the summer, resulted in lower milk performance and lactation persistence than in the IF system. The findings suggest that the PF cows were not able to exploit their genetic milk performance potential, which confirms the results of Australian HF cows with a slightly higher level of concentrate (Fulkerson *et al.* 2008). Additional feeding with energy-rich feed, e.g. coarse meal of maize ears, would have compensated but would have resulted in additional AL purchased outside the farm, which would not be consistent with a low-cost, self-sufficient PF system. Moving the cows to Alpine pasture might also be a management option to compensate such summer feed lack. Indeed, this is common practice in Alpine and pre-Alpine regions.

Economic efficiency

The higher milk price of the IF herd is explained by higher milk solids content and a more constant daily milk yield over the year. The higher milk fat quality (higher content of CLA and $\omega-3$ fatty acids) in PF milk did not yield additional profits. The market demands a constant supply of milk, which PF inherently cannot provide due to seasonality.

The higher fixed costs in the IF system resulted especially from the higher machinery and conservation costs for maize and silage. The findings of the present study are consistent with simulation studies by Gazzarin *et al.* (2005; Gazzarin & Lips 2006) and retrospective observational studies by Hofstetter (2010) and Gazzarin *et al.* (2011).

Previous calculations have concluded that PF and IF systems with the same AL appear to be on a par in terms of economic efficiency, but only if the IF cows show a considerably higher milk yield than in the present study (Gazzarin *et al.* 2005; Gazzarin & Lips 2006). A difference in milk yield of more than 2500 kg milk per cow and lactation between PF and IF do not compensate for the higher fixed costs. This is explained by high variable costs for purchased concentrate and also the contractor costs of feed production, which can hardly be reduced by economies of scale. The higher milk price does not, however, compensate for the high total costs of IF, leading to a much lower net income.

Farms with limited AL should reduce their direct and fixed costs (i.e. feed and machinery) as a fast and

targeted way to increase profits. The lower milk yield in PF cows is more than compensated for by the lower fixed costs and by a higher return in by-products such as more sales of surplus calves. The surplus calves will gain in importance due to increased demand for meat. Furthermore, a recent comprehensive, model-based study by Zehetmeier *et al.* (2012) put the specialization in both dairy and beef production in terms of greenhouse gas emissions in question. Their findings showed higher human-edible efficiency, i.e. output human-edible protein/cow to input human-edible food, for dairy cows yielding 6000 kg of milk/year, e.g. the dual-purpose Fleckvieh breed, compared with dairy-only cows yielding 10000 kg of milk/year, e.g. the HF breed. Due to the co-products they yield, dual-purpose cattle such as SF or Montbéliard might undergo a renaissance, especially for PF systems. Furthermore, higher returns from by-products will decrease the income risk of high milk price volatility.

CONCLUSIONS

The results of the present study suggest that IF cows are more efficient in terms of milk performance, milk fat and milk protein production compared with PF cows. The milk production performance and FA composition of milk from PF cows are affected by seasonal changes in the availability and nutritional content of grasses. The lower feeding intensity and the better animal welfare conditions probably resulted in PF cows showing a higher fertility than IF cows. Better commercialisation of the higher content of CLA and $\omega-3$ FA of the PF system could result in a higher milk price.

The IF herds had higher land productivity than PF herds. However, to achieve a lactation performance of about 9000 kg ECM, soyabean meal and maize gluten constituted about 0.2 of the IF AL and this feed must usually be purchased, whereas a PF farm with a lactation performance of c. 6000 kg ECM is almost self-sufficient. The PF farm also achieved better economic results due to its lower costs. Within the current situation and the associated costs in Switzerland, the more economically interesting strategy would be to lower costs rather than to increase the milk yield. Furthermore, dual-purpose cattle in PF systems could have advantages regarding its lower use of human-edible food and the increasing global demand for meat.

Rising feed prices all over the world suggest that pasture-based dairy production systems could gain importance in the foreseeable future on AL with humid

climate conditions. Pasture-land systems already dominate in countries such as Ireland or New Zealand. In an environment of rising input costs and milk price volatility, farmers are under pressure to improve their management systems and must therefore be able to accurately assess the profitability and risk impact of the system they are using. Future larger-scale and longer-term studies should supply further evidence for the economic efficiency of different production systems.

This study was funded by the Swiss Dairy Producers' Association, the Central Swiss Dairy Producers' Association, the Departments of Education & Culture and Building, Environment & Economics of the Canton of Lucerne, and AGFF Zurich. The authors also wish to thank our colleagues at our institutions and the staff at the experimental farm in Hohenrain, Switzerland, especially Josef Estermann. A part of this study was published in *Agrarforschung Schweiz* 2(9): 402–423, 2011.

REFERENCES

- AGFF (Arbeitsgemeinschaft zur Förderung des Futterbaues) (2002). *Wettbewerb Flächenproduktivität in der Milchproduktion*. Zürich, Switzerland: AGFF.
- ALP (Agroscope Liebefeld-Posieux) (2008). *Fütterungsempfehlungen und Nährwerttabellen für Wiederkäuer: das Grüne Buch*. Posieux, Switzerland: Agroscope.
- ALQAISI, O., STEGLICH, J., NDAMBI, A. & HEMME, T. (2011). Feeding systems: an assessment of dairy competitiveness. In *IFCN Dairy Report 2011* (Ed. T. Hemme), pp. 174–175. Kiel, Germany: IFCN Dairy Research Center.
- AOAC (2005). *Official Methods of Analysis*. Arlington, VA, USA: AOAC.
- BARB, C. R. & KRAELING, R. R. (2004). Role of leptin in the regulation of gonadotropin secretion in farm animals. *Animal Reproduction Science* **82**, 155–167.
- BARGO, F., MÜLLER, L. D., DELAHOY, J. E. & CASSIDY, T. W. (2002). Performance of high producing dairy cows with three different feeding systems combining pasture and total mixed rations. *Journal of Dairy Science* **85**, 2948–2963.
- BRAUNVIEH SCHWEIZ (2011). *Geschäftsbericht 2010*. Zug, Switzerland: Braunvieh Schweiz. Available from: http://homepage.braunvieh.ch/xml_1/internet/de/application/d7/d999/f1103.cfm (accessed 14 February 2014).
- BRAUNVIEH SCHWEIZ (2012). *Zuchtwertschätzung*. Zug, Switzerland: Braunvieh Schweiz. Available from: http://homepage.braunvieh.ch/xml_1/internet/de/application/d4/f58.cfm (accessed 5 February 2014).
- BRAUNVIEH SCHWEIZ (2013). *Vorschriften MLP*. Zug, Switzerland: Braunvieh Schweiz. Available from: http://homepage.braunvieh.ch/xml_1/internet/de/application/d7/d548/f550.cfm (accessed 14 January 2014).
- BURREN, A., GAZZARIN, CH., KECKEIS, K., KUNZ, P., PICCAND, V., PITT-KÄCH, S., RIEDER, ST., ROTH, N., SCHORI, F., THOMET, P., TROXLER, J., WANNER, M. & WEILENMANN, S. (2010). *Projekt Weidekuhgenetik, 2007–2010. Schlussbericht*. (Eds. N. Roth & P. Kunz), Zollikofen, Switzerland: Schweizerische Hochschule für Landwirtschaft. Available from: <http://dx.doi.org/10.5167/uzh-60075> (accessed 5 February 2014).
- BUTLER, W. R. (2005). Relationship of negative energy balance with fertility. *Advances in Dairy Technology* **17**, 35–46.
- BUTLER, W. R. & SMITH, R. D. (1989). Interrelationship between energy balance and postpartum reproductive function in dairy cattle. *Journal of Dairy Science* **72**, 767–783.
- COLEMAN, J., PIERCE, K. M., BERRY, D. P., BRENNAN, A. & HORAN, B. (2009). The influence of genetic selection and feed system on the reproductive performance of spring-calving dairy cows within future pasture-based production systems. *Journal of Dairy Science* **92**, 5258–5269.
- COLLOMB, M. & BÜHLER, T. (2000). Analyse de la composition en acides gras de la graisse de lait. I. Optimisation et validation d'une méthode générale à haute résolution. *Mitteilungen aus Lebensmitteluntersuchung und Hygiene* **91**, 306–332.
- COLLOMB, M., BISIG, W., BÜTIKOFER, U., SIEBER, R., BREGY, M. & ETTER, L. (2008). Seasonal variation in the fatty acid composition of milk supplied to dairies in the mountain regions of Switzerland. *Dairy Science and Technology* **88**, 631–647.
- CORRAL, A. J. & FENLON, J. S. (1978). A comparative method for describing the seasonal distribution of production from grasses. *Journal of Agricultural Science, Cambridge* **91**, 61–67.
- CURRAN, J., DELABY, L., KENNEDY, E., MURPHY, J. P., BOLAND, T. M. & O'DONOVAN, M. (2010). Sward characteristics, grass dry matter intake and milk production performance are affected by pre-grazing herbage mass and pasture allowance. *Livestock Science* **127**, 144–154.
- DAGET, P. & POISSONET, J. (1969). *Analyse Phytologique des Prairies. Application Agronomique*. Document no. 48. Montpellier, France: CNRS.
- DHIMAN, T. R., ARNAND, G. R., SATTER, L. D. & PARIZA, M. W. (1999). Conjugated linoleic acid content of milk from cows fed different diets. *Journal of Dairy Science* **82**, 2146–2156.
- DILLON, P., ROCHE, J. R., SHALLOO, L. & HORAN, B. (2005). Optimising financial return from grazing in temperate pastures. In *Utilisation of Grazed Grass in Temperate Animal Systems. Proceedings of a Satellite Workshop of the XXth International Grassland Congress, Cork Ireland* (Ed. J. J. Murphy), pp. 131–138. Wageningen, The Netherlands: Wageningen Academic Publishers.
- DURGIAL, B. (1996). Mit Kurzrasenweide die Milchproduktionskosten senken. *Agrarforschung* **3**, 509–512.
- EDMONSON, A. J., LEAN, I. J., WEAVER, L. D., FARVER, T. & WEBSTER, G. (1989). A body condition scoring chart for Holstein dairy cows. *Journal of Dairy Science* **72**, 68–78.

- FAO (2013). *The State of Food and Agriculture*. Rome: FAO. Available from: <http://www.fao.org/docrep/018/i3301e/i3301e.pdf> (accessed 14 January 2014).
- FARIÑA, S. R., GARCIA, S. C., FULKERSON, W. J. & BARCHIA, I. M. (2011). Pasture-based dairy farm systems increasing milk production through stocking rate or milk yield per cow: pasture and animal responses. *Grass and Forage Science* **66**, 316–332.
- FLISCH, R., SINAJ, S., CHARLES, R. & RICHNER, W. (2009). GRUDAF: Grundlagen für die Düngung im Acker- und Futterbau. *Agrarforschung* **16**, 1–100.
- FONTANELI, R. S., SOLLENBERGER, L. E., LITTELL, R. C. & STAPLES, C. R. (2005). Performance of lactating dairy cows managed on pasture-based or freestall barn-feeding systems. *Journal of Dairy Science* **88**, 1264–1276.
- FULKERSON, W. J., DAVISON, T. M., GARCIA, S. C., HOUGH, G., GODDARD, M. E., DOBOS, R. & BLOCKEY, M. (2008). Holstein–Friesian dairy cows under a prodominantly grazing system: interaction between genotype and environment. *Journal of Dairy Science* **91**, 826–839.
- GAZZARIN, CH. & HILTY, R. (2002). *Stallsysteme für Milchvieh: Vergleich der Bauinvestitionen*. FAT-Bericht no. 586. Tänikon, Switzerland: FAT.
- GAZZARIN, CH. & LIPS, M. (2006). *Entwicklungsoptionen typischer Milchproduktionsbetriebe unter AP 2011 – Investieren, Spezialisieren oder Kooperieren?* FAT-Bericht no. 651. Ettenhausen, Switzerland: Agroscope FAT Tänikon.
- GAZZARIN, CH. & LIPS, M. (2007). *Lohnt sich das Wachstum in der Milchproduktion? Dynamische Simulation für 30 Jahre*. ART-Bericht no. 693. Ettenhausen, Switzerland: Forschungsanstalt Agroscope Reckenholz-Tänikon ART.
- GAZZARIN, CH. & SCHICK, M. (2004). *Milchproduktionssysteme für die Talregion – Vergleich von Wirtschaftlichkeit und Arbeitsbelastung*. FAT-Bericht no. 608. Tänikon, Switzerland: FAT.
- GAZZARIN, CH., AMMANN, H., SCHICK, M., VAN CAENEGEM, L. & LIPS, M. (2005). *Milchproduktionssysteme in der Tal- und Hügelregion – was ist optimal für die Zukunft?* FAT-Bericht no. 645. Ettenhausen, Switzerland: Agroscope FAT Tänikon.
- GAZZARIN, CH., BRAND, R., ALBISSER, G. & WETTSTEIN, N. (2011). *Milchproduktion auf Berg- und Hügelbetrieben in der Schweiz und Österreich*. ART-Bericht no. 749. Ettenhausen, Switzerland: Forschungsanstalt Agroscope Reckenholz-Tänikon ART.
- HILTY, R., VAN CAENEGEM, L. & HERZOG, D. (2007). *ART-Preisbaukasten 2007. Baukostensammlung für Landwirtschaftliche Betriebsgebäude*. Ettenhausen, Switzerland: Forschungsanstalt Agroscope Reckenholz-Tänikon.
- HODEN, A., FIORELLEI, J. L., JEANNIN, B., HUGUET, L., MULLER, A. & WEISS, P. (1987). Le pâturage simplifié pour vaches laitières: synthèse de résultats expérimentaux. *Fourrages* **111**, 239–257.
- HOFSTETTER, P. (2010). Structural analysis of dairy industry and its evolution in Central Switzerland. *Grassland Science in Europe* **15**, 109–111.
- HOFSTETTER, P., STEIGER BURGOS, M., PETERMANN, R., MÜNGER, A., BLUM, J. W., THOMET, P., MENZI, H., KOHLER, S. & KUNZ, P. (2011). Does body size of dairy cows, at constant ratio of maintenance to production requirements, affect productivity in a pasture-based production system? *Journal of Animal Physiology and Animal Nutrition* **95**, 717–729.
- HOLDEN, L. A., MULLER, L. D. & FALES, S. L. (1994). Estimation of intake in high producing Holstein cows grazing grass pasture. *Journal of Dairy Science* **77**, 2332–2340.
- Holstein Switzerland (2014). *Gesamtindex ISEL*. Posieux, Switzerland: Holstein Switzerland. Available from: <http://www.holstein.ch/rubriques/?keyRubrique=gesamtindex> (accessed 14 February 2014).
- HORAN, B., DILLON, P., FAVERDIN, P., DELABY, L., BUCKLEY, F. & RATH, M. (2005). The interaction of strain of Holstein–Friesian cows and pasture-based feed system on milk yield, body weight and body condition score. *Journal of Dairy Science* **88**, 1231–1243.
- HORN, M., STEINWIDDER, A., GASTEINER, J., PODSTATZKY, L., HAIGER, A. & ZOLLITSCH, W. (2013). Suitability of different dairy cow types for an Alpine organic and low-input milk production system. *Livestock Science* **153**, 135–146.
- KAUFMANN, L. D., MÜNGER, A., RÉRAT, M., JUNGHANS, P., GÖRS, S., METGES, C. C. & DOHME-MEIER, F. (2011). Energy expenditure of grazing cows and cows fed grass indoors as determined by the 13 C bicarbonate dilution technique using an automatic blood sampling system. *Journal of Dairy Science* **94**, 1989–2000.
- KOLVER, E. S. & MULLER, L. D. (1998). Performance and nutrient intake of high producing Holstein cows consuming pasture or a total mixed ration. *Journal of Dairy Science* **81**, 1403–1411.
- LINN, J. (2013). *Feeding Total Mixed Rations*. Dairy Extension. St. Paul, MN, USA: University of Minnesota/Extension. Available from: <http://www1.extension.umn.edu/agriculture/dairy/feed-and-nutrition/feeding-total-mixed-rations/> (accessed 15 January 2014).
- MACDONALD, K. A., PENNO, J. W., LANCASTER, J. A. S. & ROCHE, J. R. (2008). Effect of stocking rate on pasture production, milk production, and reproduction of dairy cows in pasture-based systems. *Journal of Dairy Science* **91**, 2151–2163.
- MCÉVOY, M., KENNEDY, E., MURPHY, J. P., BOLAND, T. M., DELABY, L. & O'DONOVAN, M. (2008). The effect of herbage allowance and concentrate supplementation on milk production performance and dry matter intake of spring-calving dairy cows in early lactation. *Journal of Dairy Science* **91**, 1258–1269.
- MCÉVOY, M., O'DONOVAN, M., KENNEDY, E., MURPHY, J. P., DELABY, L. & BOLAND, T. M. (2009). Effect of pregrazing herbage mass and pasture allowance on the lactation performance of Holstein–Friesian dairy cows. *Journal of Dairy Science* **92**, 414–422.
- METZNER, M., HEUWIESER, W. & KLEE, W. (1993). Die Beurteilung der Körperkondition im Herdenmanagement (Body condition scoring in dairy herd health management). *Der Praktischer Tierarzt* **11**, 991–998.
- MOSIMANN, E. (2001). Croissance des herbages. Méthodes de mesures et applications pratiques. *Revue Suisse Agriculture* **33**, 163–167.

- MÜNGER, A. & JANS, F. (2001). Umtriebs- und Kurzrasenweide für Milchkühe im Vergleich. *Agrarforschung Schweiz* **8**, 464–469.
- NRC (2001). *Nutrient Requirements of Dairy Cattle*, 7th edn. (revised edition). Washington, DC: National Academy Press.
- NYFELER, D., HUGUENIN-ELIE, O., SUTER, M., FROSSARD, E. & LÜSCHER, A. (2011). Grass–legume mixtures can yield more nitrogen than legume pure stands due to mutual stimulation of nitrogen uptake from symbiotic and non-symbiotic sources. *Agriculture, Ecosystems and Environment* **140**, 155–163.
- PULIDO, R.G. & LEAVER, J.D. (2003). Continuous and rotational grazing of dairy cows – the interactions of grazing system with level of milk yield, sward height and concentrate level. *Grass and Forage Science* **58**, 265–275.
- R Development Core Team (2012). *R Software Version 2.11.1*. Available from: <http://www.r-project.org/> (accessed 15 January 2014).
- REIST, M., KOLLER, A., BUSATO, A., KUPFER, U. & BLUM, J.W. (2000). First ovulation and ketone body status in the early postpartum period of dairy cows. *Theriogenology* **54**, 685–701.
- SCHORI, F. (2009). Weidebesatzstärken: Auswirkung auf Milchleistung und Grasqualität. *Agrarforschung Schweiz* **16**, 436–441.
- SCHORI, F. & MÜNGER, A. (2010). Grazing behavior and intake of two Holstein cow types in a pasture-based production system. *Grassland Science in Europe* **15**, 895–897.
- SJAUNJA, L.O., BAEVRE, L., JUNKKARINEN, L., PEDERSEN, J. & SETÄLÄ, J. (1990). A Nordic proposal for an energy corrected milk (ECM) formula. In *Performance Recording of Animals: State of the Art, 1990. Proceedings of the 27th Biennial Session of the International Committee for Animal Recording (ICAR), July 2–6 1990, Paris, France*. EEAP Publication no. 50 (Ed. J.A.M. van Arendonk), pp. 156–157. Wageningen, The Netherlands: Pudoc.
- STARK, R., SCHICK, M. & MORIZ, C. (2009). *ART-Arbeitsvoranschlag. Planungsinstrument zur Kalkulation des Personal und Maschineneinsatzes auf landwirtschaftlichen Betrieben. Software Version 2009*. Ettenhausen, Switzerland: Forschungsanstalt Agroscope Reckenholz-Tänikon ART.
- STEIGER BURGOS, M., PETERMANN, R., HOFSTETTER, P., THOMET, P., KOHLER, S., MÜNGER, A., BLUM, J.W., MENZI, H. & KUNZ, P. (2007). Quel type de vache laitière pour produire du lait au pâturage? *Revue Suisse Agriculture* **39**, 123–128.
- Swiss Herdbook (2013). *Reglement über die Zuchtwertschätzung und Nachzuchtprüfung bei swissherdbook*. Zollikofen, Switzerland: Swiss Herdbook. Available from: <http://www.swissherdbook.ch/genetik/zuchtwerte/> (accessed 5 February 2014).
- THOMET, P. & BLÄTTLER, T. (1998). Graswachstum als Grundlage für die Weideplanung. *Agrarforschung* **5**, 25–28.
- THOMET, P., HADORN, M., JANS, F., TROXLER, J., PERLER, O. & MEILI, E. (1999). *Kurzrasenweide Intensivstandweide. Arbeitsgemeinschaft zur Förderung des Futterbaus (AGFF) Merkblatt, 2*. Zurich, Switzerland: FAL Zürich-Reckenholz.
- THOMET, P., PICCAND, V., SCHORI, F., TROXLER, J., WANNER, M. & KUNZ, P. (2010). Efficiency of Swiss and New Zealand dairy breeds under grazing conditions on Swiss dairy farms. *Grassland Science in Europe* **15**, 1018–1020.
- TOZER, P.R., BARGO, F. & MULLER, L.D. (2003). Economic analyses of feeding systems combining pasture and total mixed ration. *Journal of Dairy Science* **86**, 808–818.
- VANCE, E.R., FERRIS, C.P., ELLIOTT, C.T., MCGETRICK, S.A. & KILPATRICK, D.J. (2012). Food intake, milk production, and tissue changes of Holstein–Friesian and Jersey × Holstein–Friesian dairy cows within a medium-input grazing system and a high-input total confinement system. *Journal of Dairy Science* **95**, 1527–1544.
- VAN SOEST, P.J., ROBERTSON, J.B. & LEWIS, B.A. (1991). Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science* **74**, 3583–3597.
- VIBART, R.E., FELLNER, V., BURNS, J.C., HUNTINGTON, G.B. & GREEN, J.T. Jr (2008). Performance of lactating dairy cows fed varying levels of total mixed ration and pasture. *Journal of Dairy Research* **75**, 471–480.
- VIBART, R.E., WASHBURN, S.P., GREEN, J.T. Jr, BENSON, G.A., WILLIAMS, C.M., PACHECO, D. & LOPEZ-VILLALOBOS, N. (2012). Effect of feeding strategy on milk production, reproduction, pasture utilization, and economics of autumn-calving dairy cows in eastern North Carolina. *Journal of Dairy Science* **95**, 997–1010.
- WexTech Systems, Inc. (2006). *CPM-Dairy Version 3.0.8*. Kennett Square, PA, USA/Ithaca, NY, USA/Chazy, NY, USA: The Center for Animal Health and Productivity, School of Veterinary Medicine, University of Pennsylvania/The Department of Animal Science, Cornell University/The William H. Miner Agricultural Research Institute.
- WHITE, S.L., BERTRAND, J.A., WADE, M.R., WASHBURN, S.P., GREEN, J.T. & JENKINS, T.V. (2001). Comparison of fatty acid content of milk from Jersey and Holstein cows consuming pasture or total mixed ration. *Journal of Dairy Science* **84**, 2295–2301.
- WYSS, U., MAURER, J., FREY, HJ., REINHARD, TH., BERNET, A. & HOFSTETTER, P. (2011). Systemvergleich Milchproduktion Hohenrain. Aspekte der Milchqualität und der Saisonalität der Milcheinlieferungen. *Agrarforschung Schweiz* **2**, 412–417.
- ZEHETMEIER, M., BAUDRACCO, J., HOFFMANN, H. & HEISSENHUBER, A. (2012). Dose increasing milk yield per cow reduce greenhouse gas emissions? A system approach. *Animal* **6**, 154–166.
- ZIMMERMANN, A. (2006). *Kosten und Umweltwirkung der Milchviehfütterung*. ART-Bericht no. 662. Ettenhausen, Switzerland: Forschungsanstalt Agroscope Reckenholz-Tänikon ART.